

Our Ref. No.: 002013.P002
Express Mail No. EL651893343US

UTILITY APPLICATION FOR UNITED STATES PATENT

FOR

BISPHENYLENE-SPIROBIFLUORENE COMPOUNDS, METHOD FOR SYNTHESIZING
THE SAME, AND ELECTROLUMINESCENCE MATERIAL AND DEVICE HAVING THE
SAME

Inventor(s): Hyoyoung Lee et al.

327253
 327254
 327255
 327256
 327257
 327258
 327259
 327260
 327261
 327262
 327263
 327264
 327265
 327266
 327267
 327268
 327269
 327270
 327271
 327272
 327273
 327274
 327275
 327276
 327277
 327278
 327279
 327280
 327281
 327282
 327283
 327284
 327285
 327286
 327287
 327288
 327289
 327290
 327291
 327292
 327293
 327294
 327295
 327296
 327297
 327298
 327299
 327300
 327301
 327302
 327303
 327304
 327305
 327306
 327307
 327308
 327309
 327310
 327311
 327312
 327313
 327314
 327315
 327316
 327317
 327318
 327319
 327320
 327321
 327322
 327323
 327324
 327325
 327326
 327327
 327328
 327329
 327330
 327331
 327332
 327333
 327334
 327335
 327336
 327337
 327338
 327339
 327340
 327341
 327342
 327343
 327344
 327345
 327346
 327347
 327348
 327349
 327350
 327351
 327352
 327353
 327354
 327355
 327356
 327357
 327358
 327359
 327360
 327361
 327362
 327363
 327364
 327365
 327366
 327367
 327368
 327369
 327370
 327371
 327372
 327373
 327374
 327375
 327376
 327377
 327378
 327379
 327380
 327381
 327382
 327383
 327384
 327385
 327386
 327387
 327388
 327389
 327390
 327391
 327392
 327393
 327394
 327395
 327396
 327397
 327398
 327399
 327400
 327401
 327402
 327403
 327404
 327405
 327406
 327407
 327408
 327409
 327410
 327411
 327412
 327413
 327414
 327415
 327416
 327417
 327418
 327419
 327420
 327421
 327422
 327423
 327424
 327425
 327426
 327427
 327428
 327429
 327430
 327431
 327432
 327433
 327434
 327435
 327436
 327437
 327438
 327439
 327440
 327441
 327442
 327443
 327444
 327445
 327446
 327447
 327448
 327449
 327450
 327451
 327452
 327453
 327454
 327455
 327456
 327457
 327458
 327459
 327460
 327461
 327462
 327463
 327464
 327465
 327466
 327467
 327468
 327469
 327470
 327471
 327472
 327473
 327474
 327475
 327476
 327477
 327478
 327479
 327480
 327481
 327482
 327483
 327484
 327485
 327486
 327487
 327488
 327489
 327490
 327491
 327492
 327493
 327494
 327495
 327496
 327497
 327498
 327499
 327500
 327501
 327502
 327503
 327504
 327505
 327506
 327507
 327508
 327509
 327510
 327511
 327512
 327513
 327514
 327515
 327516
 327517
 327518
 327519
 327520
 327521
 327522
 327523
 327524
 327525
 327526
 327527
 327528
 327529
 327530
 327531
 327532
 327533
 327534
 327535
 327536
 327537
 327538
 327539
 327540
 327541
 327542
 327543
 327544
 327545
 327546
 327547
 327548
 327549
 327550
 327551
 327552
 327553
 327554
 327555
 327556
 327557
 327558
 327559
 327560
 327561
 327562
 327563
 327564
 327565
 327566
 327567
 327568
 327569
 327570
 327571
 327572
 327573
 327574
 327575
 327576
 327577
 327578
 327579
 327580
 327581
 327582
 327583
 327584
 327585
 327586
 327587
 327588
 327589
 327590
 327591
 327592
 327593
 327594
 327595
 327596
 327597
 327598
 327599
 327600
 327601
 327602
 327603
 327604
 327605
 327606
 327607
 327608
 327609
 327610
 327611
 327612
 327613
 327614
 327615
 327616
 327617
 327618
 327619
 327620
 327621
 327622
 327623
 327624

BISPHENYLENE-SPIROBIFLUORENE COMPOUNDS, METHOD FOR
SYNTHESIZING THE SAME, AND ELECTROLUMINESCENCE MATERIAL AND
DEVICE HAVING THE SAME

5

BACKGROUND OF THE INVENTION

1. Field of the Invention

10 The present invention relates to an organic electroluminescence (EL) compound, a method for synthesizing the same, and an EL material and device having the same, and more particularly, to bisphenylene-spirobifluorene compounds, a method for synthesizing the same, and EL material and device having the same.

2. Description of the Related Art

15 In general, an electroluminescence (EL) device using low molecular weight materials is formed by sequentially depositing a transparent electrode, an electrically conducting buffer layer, a light-emitting layer, an electron transport layer and a metal electrode on a substrate under high vacuum. If the transparent electrode and the metal electrode are connected to positive (+) and negative (-) terminals of power, holes are supplied to the hole transport layer and the light-emitting layer through the transparent electrode and electrons are supplied to the light-emitting layer through the metal electrode. The holes and electrons supplied to the light-emitting layer are combined therein, thereby achieving luminescence. Since organic EL devices are quick in response, driven at a low operating voltage and self-illuminating, they require no additional back light source, making them small and lightweight. Also, the organic EL devices have excellent luminance and are not dependent upon viewing angle of display.

20

25

Types of light-emitting materials forming the organic EL devices include low molecular weight organic materials and organic polymer materials. While the low molecular weight organic material is subjected to vacuum deposition to form a thin film on a substrate, the organic polymer material is molten in an organic solvent to be subjected to spin coating to form a thin film on a substrate. The low molecular

30

weight organic luminescence device is described in detail in U.S. Patent No. 4,539,507.

Since the color of light emitted is varied according to the band gap of light-emitting molecule, it is quite important to develop red (R), green (G) and blue (B) emission materials having high luminance at low voltages while maintaining high color purity according to the wavelength of light emitted. Development of B emission materials is, inter alia, highly demanded. This is because one of full-color organic EL device implementation methods, that is, RGB emission, can be implemented by forming a blue emission layer, depositing a color change material (CCM) capable of adjusting wavelengths and converting blue emission having the highest energy into green and red emissions.

As materials to emit blue fluorescent light, polyphenylenes having phenylene as a basic unit were reported by J. M. Tour in 1991 (Adv. Mater. 1994, 6, No. 3, pp. 190-198). However, these materials have poor solubility while they are excellent in view of thermal stability and color purity. Also, since it was known that materials having fluorenes, which are pseudo materials of phenylenes, as a basic unit, are promising, a great deal of research into these materials has been made hereto.

Woo, Inbasekaran, Shiang and Roof developed polymer EL polymers using a polymerization technique of polymers, called Suzuki coupling, and disclosed poly(fluorene) based EL materials in WO 97/05184 in 1997. In 1999, poly(fluorenes) with polar side chains were proposed by Pei, Yu and Yang, but the polar side chains promote formation of excimer, which is one of problems with poly(fluorene), resulting in degradation in color purity (see U.S. Patent No. 5,900,327 and J. Appl. Phys., Vol. 81, pp 3294-3298). In order to solve the problem of degradation in color purity of blue light-emitting electrochemical cells due to formation of excitons by conventional poly(fluorenes), Chen, Klaerner, Miller and Scott of IBM Corporation developed poly(anthracene-co-fluorene) copolymers obtained by copolymerizing the poly(fluorenes) with anthracenes (see U.S. Patent No. 5,998,045). As described above, fluorene-based B light-emitting materials with high blue color purity and emission efficiency have been developed. However,

planarity of fluorene molecules causes an excimer phenomenon in which the fluorene molecules get easily closer to each other than non-planar molecules, resulting in a change in wavelength of light emitted. The excimer phenomenon makes it difficult to maintain color purity.

5 In practice, one of B emission materials being currently in widespread use is a compound of distyrylarylene (DSA) based 1,4-bis(2,2-diphenylvinyl)biphenyl (DPVBi) proposed by Hosokawa, Higashi, Nakamura and Kusumoto, in Appl. Phys. Lett. 67 (26), 3853, 1995, and manufactured by Idemitsu Kosan Co., Ltd. Here, the luminance was 6,000 cd/m² and the emission efficiency was 0.7 to 0.8 lm/W. Also, it was reported that the DSA-based compound improved approximately two times in luminance and emission efficiency, that is, 10,000 cd/m² and 1.5 lm/W, when it is used with a doping material. The DPVBi used herein has a non-planar molecular structure due to steric inhibition of an arylene substituent.

10 Accordingly, preparation of spirobifluorene compounds having properties of fluorenes and non-planar molecular structures, and organic EL devices thereof, are being vigorously studied.

15 Kreuder, Lupo, Salbeck, Schenk and Stehlin disclosed poly(spirobifluorenes) having a spiro framework and copolymers thereof (see U.S. Patent No. 5,621,131). Also, they disclosed EL materials of new spiro low molecular weight materials or polymers having a spiro framework other than spirobifluorene or having a hetero element substituted for 9,9'-positioned carbon element (see U. S. Patent Nos. 20 5,741,921, 5,763,636 and 5,859,211). U.S. Patent No. 5,840,271 to Lupo et al. discloses methods of synthesizing low molecular weight and polymer compounds of 2,7-diphenylene-9,9'-spirobifluorene and 2,2'-diphenylene-9,9'-spirobifluorene having 25 two identical phenylene groups substituted in 2,7- or 2,2'- positions. Although the structure free of alkyl substituent was effective in maintaining color purity and color stability, the processibility of the low molecular weight or polymer materials was poor. In other words, since such material has poor solubility in organic solvents, its film-forming properties are poor.

In order to overcome the poor film-forming properties, Rietz et al. reported in U.S. Patent No. 6,132,641 that 2,2'-dialkyl-9,9'-bisspirofluorene having alkyl side chains introduced in 2,2'- positions and its polymer 2,2'-dialkyl-7,7'-poly(9,9'-spirobifluorene) have improved solubility in organic solvents and improved polymer film forming properties in EL device applications.

SUMMARY OF THE INVENTION


To solve the above-described problems, it is a first object of the present invention to provide alkyl substituted bisphenylene-spirobifluorene compounds configured to improve film forming properties while maintaining high blue color purity and enhancing color stability of spirobifluorene.

It is a second object of the present invention to provide a method of preparing bisphenylene-spirobifluorene compounds, based on spirobifluorene molecular structure, which can substitute alkyl side chains at desired positions to increase film forming properties.

It is a third object of the present invention to provide EL materials containing new bisphenylene-spirobifluorene compounds.

It is a fourth object of the present invention to provide an EL device having a light-emitting layer containing a new bisphenylene-spirobifluorene compound having a structure in which alkyl substituent is introduced at a new position to implement a device with high efficiency, high color purity and excellent color stability.

To accomplish the first object, the present invention provides a bisphenylene-spirobifluorene compound defined by the following formula:



Wherein R_1 and R_2 are identical or different and are independently a straight-chain or branched alkyl group having from 1 to 22 carbon atoms, X_1 and X_2 independently contains one or more elements selected from the group consisting of C, O, N, S, Si and Ge, and m and n are integers from 1 to 4. Preferably, X_1R_1 and X_2R_2 are at 1',6'-positions.

To achieve the second object of the present invention, a method of preparing bisphenylene-spirobifluorene compound includes the step of forming a biphenyl compound having X_1R_1 and X_2R_2 in which R_1 and R_2 are identical or different and are independently a straight-chain or branched alkyl group having from 1 to 22 carbon atoms, and X_1 and X_2 independently contains one or more elements selected from the group consisting of C, O, N, S, Si and Ge. Then, a biphenyl-halogenated fluoreneol compound having the X_1R_1 and X_2R_2 is formed by reacting the biphenyl compound with halogenated fluorenone. A halogenated spirobifluorene compound having the X_1R_1 and X_2R_2 is formed from the biphenyl-halogenated fluoreneol compound by cyclization. Then, a bisphenylene-spirobifluorene compound having the X_1R_1 and X_2R_2 is formed by substituting halogen of the halogenated spirobifluorene compound by a phenyl group.

In the step of forming the biphenyl-halogenated fluoreneol compound, a metal-halogen ligand substitution reaction is preferably employed.

Also, in the step of forming the biphenyl-halogenated fluoreneol compound, a slightly excess amount of halogenated fluorene-one is preferably added.

To achieve the third object of the present invention, there is provided an electroluminescence (EL) material comprising the bisphenylene-spirobifluorene compound defined by the formula stated above. The bisphenylene-spirobifluorene compound is preferably contained in an amount of 10% by weight or more.

To achieve the fourth object of the present invention, there is provided an electroluminescence (EL) device including a cathode, an anode, and a light-emitting layer interposed between the cathode and the anode and containing the EL material according to the present invention. Preferably, the bisphenylene-spirobifluorene compound is contained in the light emitting layer in an amount of 10% to 100% by weight.

The bisphenylene-spirobifluorene compound according to the present invention has a structure in which alkyl substituent is introduced at a new position to increase the solubility in organic solvents and reduce excimer formation. Its color purity and color stability are good and it is highly soluble in organic solvents, thereby providing excellent processibility. Also, the bisphenylene-spirobifluorene compound according to the present invention can provide blue organic EL materials having excellent properties as electronic materials, including light transmissivity, environment resistance, adhesive strength with respect to a substrate, film forming properties, electrical field stability. Further, it is possible to remarkably reduce crystallization, which frequently occurs to a planar organic matter during formation of a thin film in forming a light-emitting layer of an EL device, thereby prolonging the life cycle of the EL device.

BRIEF DESCRIPTION OF THE DRAWINGS

The above objects and advantages of the present invention will become more apparent by describing in detailed preferred embodiments thereof with reference to the attached drawings in which:

FIG. 1 is a proton nuclear magnetic resonance (NMR) of a compound VI-1 ($m=1$, $n=1$) prepared in Example 6;

FIG. 2 is a carbon NMR of a compound VI-1 ($m=1$, $n=1$) prepared in Example

6;

FIG. 3 is a proton NMR of a compound VI-1 ($m=2$, $n=2$) prepared in Example

7;

FIG. 4 is a carbon NMR of a compound VI-1 ($m=2$, $n=2$) prepared in Example

7;

FIG. 5 is a mass spectrum of a compound VI-1 ($m=2$, $n=2$) prepared in Example 7;

FIG. 6 is a mass spectrum of a compound prepared in Example 8;

FIG. 7 is a proton NMR of a compound VI-1 ($m=3$, $n=3$) prepared in Example

9;

FIG. 8 is a carbon NMR of a compound VI-1 ($m=3$, $n=3$) prepared in Example

7;


FIG. 9 is a cross-sectional view illustrating a method of manufacturing an electroluminescence (EL) device according to the present invention; and

FIG. 10 is a graph depicting the photoluminescence (PL) and EL spectra for an EL device manufactured in Example 10.

DETAILED DESCRIPTION OF THE INVENTION

Bisphenylene-spirobifluorene compounds according to the present invention can be prepared by a synthesis method represented by the reaction scheme 1:

[Reaction scheme 1]




wherein R_1 and R_2 are identical or different and are independently a straight-chain or branched C_1 - C_{22} alkyl group, X_1 and X_2 each contains one or more elements selected from the group consisting of C, O, N, S, Si and Ge, Y_1 and Y_2 are independently halogens, and m and n are integers from 1 to 4.

The reaction scheme 2 illustrates synthesis of a bisphenylene-spirobifluorene compound in the case where X_1 and X_2 are oxygen atom (O) in the reaction scheme 1, depicting that alkyl groups are selectively introduced into 3',6'-positions or

1',6'-positions of the spirobifluorene compound and phenyl groups are introduced into 2,7-positions.

[Reaction scheme 2]



The method of synthesizing the bisphenylene-spirobifluorene compound according to the present invention represented by the reaction scheme 2 will now be described in detail. In order to prepare a spiro compound, a halogen element must be fixed at the 2- position of biphenyl. Several different 2-bromobiphenyl derivatives having one or two alkoxy substituents are disclosed by Sparling et al. in Biomed. Mass Spectrum (1980), 7(1), 13-19.

In the reaction scheme 2, 2-halogenated dialkoxybiphenyl is used as a starting material for synthesizing a compound VI-1 of the present invention and its geometric isomer compound VI-2. First, the compound I can be prepared by mono-halogenizing dialkoxybiphenyl, and a preparation method thereof is described by Jandrue et al. in WO 9304403 and Geiger et al. in Chim. Ther. (1966), (7), 425-37.

In order to adjust the length of an alkyl chain, two alkoxy groups are substituted by two hydroxy groups to obtain the compound II, and a similar reaction is disclosed by McOmie et al. in Tetrahedron, 1968, 24, 2289. Then, the compound II having two hydroxy groups is reacted with alkyl halogen having a different chain length to obtain the compound III having C₂, C₆, C₈, C₁₂ or C₂₂ alkyl groups.

To the compound III is added 2,7-dihalogenated fluorene-9-one to obtain the compound IV. The present inventors obtained the compound IV using a metal-ligand substitution reaction. That is to say, metallic lithium and halogen ligand in the compound III were subjected to a metal-ligand substitution reaction and 2,7-dibromo fluorene-9-one was then added to obtain the compound IV. Here, in order to prevent an exchange reaction between bromine in 2,7-dibromo fluorene-9-one and metallic lithium, excess 2,7-dibromo fluorene-9-one is preferably added.

Then, cyclization of the compound IV gave the compounds V-1 and V-2 which are geometrically isomers. The cyclization was carried out by agitating the compound IV with polyphosphate at a temperature of about 50°C to about 60°C. The cyclization occurs because electrons can be easily released at 2,4-positions of an alkoxy group. In particular, the compound V-1 produced at the 4-position of an


alkoxy group at which electrons could be more easily released than at the 2-position, was produced at a higher yield when the reaction temperature was lowered.

However, under the above conditions, the yield of the compound V-1 relative to the compound V-2 was 70% or less. As the result of research conducted for finding out

more efficient reaction conditions, 94% or more of the compound V-1 relative to the compound V-2 was acquired in the presence of hydrochloric acid and acetic acid.

Finally, Suzuki reactions of the compounds V-1 and V2 with phenyl boric acid and bisphenyl boric acid provided target compounds VI-1 (in the cases where $m=n=1$ and $m=n=2$) and VI-2 (in the cases where $m=n=1$ and $m=n=2$), respectively. The

compounds VI-1 and VI-2 in which $m=n=3$, were acquired by the Suzuki reaction represented by the reaction scheme 3:



As shown in the reaction scheme 3, the compounds VI-1 and VI-2 in which $m=n=3$ were produced by Suzuki reactions of the compounds V-1 and V-2 with iodo terphenyl, respectively. The iodo terphenyl used herein was prepared by a Suzuki reaction between diiodo phenyl and phenyl boric acid.

Since the dialkyl bisphenylene-spirobifluorene derivative compound according to the present invention is a monomer having an alkyl substituent introduced to a new position, it has high solubility in organic solvents as well as excellent color purity and color stability, thereby having high processibility. Also, the present invention can provide an organic blue EL material having properties good enough to meet requirements as electronic materials, including transmissivity, environment

resistance, adhesive strength with respect to a substrate, film-forming property, electric field stability and so on.

Now, detailed examples of preparing bisphenylene-spirobifluorene compounds according to the present invention and manufacturing EL devices using the same will be described.

Example 1

Preparation of 2-bromo-5,3'-dimethoxybiphenyl

3,3'-dimethoxybiphenyl (5.0 g, 23.34 mmol) was dissolved in a dimethylformamide solvent and cooled to 0°C. N-bromosuccinimide (4.15 g, 23.34 mmol) was dissolved in a dimethylformamide solvent (63 ml) and added slowly to a solution of 3,3'-dimethoxybiphenyl for about 1.5 hours. The mixture was stirred at room temperature for about 12 hours and distilled water was added thereto, followed by stirring for about 10 minutes. The resultant was filtered three times with hexane and the filtrate was washed three times with distilled water and dried over sodium sulfate. Hexane was removed under reduced pressure to obtain a desired product (Yield: 6.70 g, 98%). The obtained product was identified as 2-bromo-5,3'-dimethoxybiphenyl through the nuclear magnetic resonance (NMR) spectrum, corresponding to the compound I of the reaction scheme 2.

δ_H (CDCl₃, 400 MHz), 7.52 (d, 1H), 7.33 (t, 1H), 6.97 (d, 1H), 6.93 (m, 2H), 6.88 (s, 1H), 6.75 (dd, 1H), 3.83 (s, 3H), 3.79 (s, 3H); δ_C (CDCl₃, 75 MHz), δ 159.1, 158.7, 143.2, 142.4, 133.7, 129.0, 121.7, 116.5, 115.0, 114.8, 113.3, 112.9, 55.5, 55.3.

Example 2

Preparation of 6-bromo-biphenyl-3,3'-diol

A solution of boron tribromide (137 ml, 1M, 137.12 mmol dissolved in dichloromethane) was slowly added to the compound I (6.70 g, 22.85 mmol) prepared in Example 1 at 0°C, followed by stirring for 1 hour. The stirred solution was allowed to warm to room temperature and continuously stirred for about 12 hours. A small amount of ice water was slowly added to the reactant mixture, followed by filtering three times with dichloromethane, washing three times with

distilled water, drying with sodium sulfate. Then, the dichloromethane solvent was removed under reduced pressure to obtain a residual product. Purification by silica gel chromatography using 4% ethyl acetate in dichloromethane gave 6-bromo-biphenyl-3,3'-diol corresponding to the compound II of the reaction scheme 2 (Yield: 5.92 g, 98%). The product was identified through the NMR spectrum that it had the identical structure with the compound II of the reaction scheme 2.

δ_H (CDCl₃, 400 MHz), 7.44 (d, 1H), 7.25 (t, 1H), 6.92 (d, 1H), 6.84 (d, 1H), 6.83 (s, 1H), 6.78 (d, 1H), 6.68 (dd, H), 5.73 (s, 1H), 5.54 (s, 1H); δ_C (CDCl₃, 75 MHz), δ 155.0, 154.9, 143.1, 142.3, 133.9, 129.2, 121.8, 118.1, 116.3, 116.2, 114.7, 112.8.

Example 3

Preparation of 2-bromo-5,3'-bis-octyloxy-biphenyl

6-bromo-biphenyl-3,3'-diol (5.92 g, 22.43 mmol) prepared in Example 2, 1-octanebromide, potassium carbonate (18.60 g, 134.56 mmol) and potassium iodide (3.72 g, 22.43 mmol) were dissolved in dimethylformamide (60 ml), followed by heating for about 24 hours at about 150°C. After cooling to room temperature, distilled water was added to the resultant and stirred for about 10 minutes, followed by filtering three times with hexane, washing three times with distilled water, drying with sodium sulfate. Then, the hexane was concentrated under reduced pressure. Purification by silica gel chromatography using 4% dichloromethane in hexane gave 2-bromo-5,3'-bis-octyloxy-biphenyl corresponding to the compound III of the reaction scheme 2 (Yield: 6.36 g, 58.1%), R₁ and R₂ in the formula of the compound III being each independently octyl. The product was identified through the NMR spectrum and element analysis that it had the identical structure with the compound III of the reaction scheme 2.

δ_H (CDCl₃, 300 MHz), 7.49 (d, 1H), 7.30 (t, 1H), 6.96-6.76 (m, 4H), 6.74 (dd, 1H), 3.98 (t, 2H), 3.91 (t, 2H), 1.80-1.74 (dt, 4H), 1.44 (dt, 4H), 1.27 (m, 16H), 0.88 (t, 6H); δ_C (CDCl₃, 75 MHz), δ 158.7, 158.3, 143.3, 142.4, 133.6, 128.9, 121.5, 117.5, 115.5, 115.3, 113.9, 112.7, 68.3, 68.1, 31.81, 31.78, 29.36, 29.30, 29.29, 29.23,

29.21, 29.17, 26.06, 25.98, 22.64, 14.1. Calc. for $C_{28}H_{41}BrO_2$: C, 68.70; H, 8.44.
Found: C, 68.70; H, 8.50.

Example 4

Preparation of 9-(5,3'-bis-octyloxy-biphenyl-2-yl)-2,7-dibromo-9H-fluorene-9-ol

5 2-bromo-5,3'-bis-octyloxy-biphenyl (0.598 g, 1.225 mmol) prepared in
Example 3 was dissolved in dried THF (30 ml) and cooled to -78°C . Then,
t-butyllithium (t-BuLi) (1.7M, 1.44 ml, 2.45 mmol) was added slowly. The obtained
resultant was stirred for about 1 hour at -78°C . After 1 hour,
10 2,7-dibromofluorene-9-one (0.497 g, 1.47 mmol) dissolved in dried THF (20 ml) was
added slowly, stirred for about 1 hour, allowed to warm to room temperature and
further stirred at room temperature for about 30 minutes. A small amount of distilled
water was added to complete the reaction and saturated brine was added to the
resultant. The organic layer was filtered twice with ether and dried over sulfate, and
the ether was concentrated under reduced pressure. Purification by silica gel
15 chromatography using 30 to 40% dichloromethane in hexane gave
9-(5,4'-bis-octyloxy-biphenyl-2-yl)-2,7-dibromo-9H-fluorene-9-ol corresponding to the
compound IV of the reaction scheme 2 (Yield: 0.62 g, 68%), R_1 and R_2 in the formula
of the compound IV being each independently octyl and Y_1 and Y_2 being bromine.
The product was identified through the NMR spectrum and element analysis that it
20 had the identical structure with the compound IV.

δ_H (CD_2Cl_2 , 400 MHz), 8.25 (d, 1H), 7.36-7.29 (m, 4H), 7.07-7.02 (m, 3H),
6.54 (t, 1H), 6.48 (d, 1H), 6.39 (dd, 1H), 5.65 (t, 1H), 5.61 (d, 1H), 3.94 (t, 2H), 3.52 (t,
2H), 2.41 (s, 1H), 1.76 (m, 2H), 1.59 (m, 2H), 1.42 (m, 4H), 1.34-1.27 (m, 16H), 0.90
(t, 3H), 0.88 (t, 3H). Calc. for $C_{41}H_{48}Br_2O_3$: C, 65.78; H, 6.46. Found: C, 65.90; H,
25 6.50.

Example 5

Preparation of 2,7-dibromo-3',6'-bisoctyloxy-9,9'-spirobisfluorene and 2,7-dibromo-1',6'-bisoctyloxy-9,9'-spirobisfluorene

The compound prepared in Example 4 (0.5 g, 0.668 mmol) was mixed with
30 acetic acid (50 ml) and hydrochloric acid (1 ml) and stirred for about 22 hours at

room temperature. After completing the reaction by adding a small amount of distilled water, the organic layer was filtered three times with dichloromethane, dried with sodium sulfate. Then, the dichloromethane was concentrated under reduced pressure. Purification by silica gel chromatography using 4% dichloromethane in hexane gave 2,7-dibromo-3',6'-bis(octyloxy)-9,9'-spirobisfluorene corresponding to the compound V-1 of the reaction scheme 2 (Yield: 0.46 g, 94%), R_1 and R_2 in the formula of the compound V-1 being each independently octyl and Y_1 and Y_2 being bromine. In the same manner, 2,7-dibromo-1',6'-bis(octyloxy)-9,9'-spirobisfluorene, which is a geometric isomer of the compound V-1, corresponding to the compound V-2 of the reaction scheme 2 (Yield: 0.024 g, 5%), R_1 and R_2 in the formula of the compound V-2 being each independently octyl and Y_1 and Y_2 being bromine. The compound V-1 of the reaction scheme 2 was prepared by recrystallization using a mixed solvent of hexane and dichloromethane. The products were identified through the NMR spectrum and element analysis that they had the identical structures with the compounds V-1 and V-2.

Compound V-1: δ_H ($CDCl_3$, 400 MHz), 7.60 (d, 2H), 7.44 (dd, 2H), 7.29 (d, 2H), 6.82 (d, 2H), 6.66 (dd, 2H), 6.56 (d, 2H), 4.02 (t, 4H), 1.80 (m, 4H), 1.47 (m, 4H), 1.31 (m, 16H), 0.88 (t, 6H), δ_C ($CDCl_3$, 400 MHz), 159.7, 151.1, 143.0, 139.6, 139.4, 130.9, 127.3, 124.6, 121.8, 121.2, 114.7, 105.9, 68.3, 64.4, 31.8, 29.4, 29.3, 26.1, 22.7, 14.1. Calc. for $C_{41}H_{46}Br_2O_2$: C, 67.40; H, 6.35. Found: C, 67.45; H, 6.40.

Compound V-2: δ_C ($CDCl_3$, 400 MHz), 7.59 (d, 2H), 7.43-7.30 (m, 5H), 6.83 (d, 2H), 6.64-6.62 (m, 2H), 6.51 (d, 1H), 4.01 (t, 2H), 3.51 (t, 2H), 1.81 (quint, 2H), 1.49 (m, 2H), 1.34-1.24 (m, 12H), 1.08 (m, 4H), 1.02 (m, 4H), 0.88 (m, 6H), δ_C ($CDCl_3$, 400 MHz), 159.6, 155.2, 150.0, 143.5, 143.0, 140.2, 139.3, 134.2, 130.4, 123.0, 126.6, 124.4, 121.4, 120.8, 114.6, 112.1, 111.1, 106.2, 68.3, 67.3, 63.7, 31.9, 31.8, 29.4, 29.3, 29.2, 28.9, 26.1, 25.4, 22.7, 22.7, 14.2, 14.1.

Example 6

Preparation of 2,7-bis(phenyl)-3',6'-bis(octyloxy)-9,9'-spirobisfluorene

The compound was prepared under Suzuki reaction conditions. The compound V-1 prepared in Example 5 (0.5 g, 0.68 mmol), 4-phenyl boric acid (0.21 g,

1.71 mmol), tetrakis(triphenylphosphine)palladium (0.24 g, 0.20 mmol), 2M-sodium carbonate (2.1 ml, 4.10 mmol) and toluene (29 ml) were simultaneously mixed, freezed under liquid nitrogen, evacuated by a vacuum pump and then warmed to room temperature to be melted. This procedure was repeated twice, followed by injection of nitrogen. The mixed solution was refluxed for 2 days and cooled to room temperature. Thereafter, a small amount of distilled water was added, and the organic layer was filtered three times with dichloromethane, dried with sodium sulfate. Then, the dichloromethane was concentrated under reduced pressure. Purification by silica gel chromatography using 10% dichloromethane in hexane, followed by recrystallization using a mixed solvent of hexane and dichloromethane gave the compound VI-1 of the reaction scheme 2 (Yield: 0.407 g, 82%), R_1 and R_2 in the formula of the compound VI-1 being each independently octyl and m and n being 1. FIGS. 1 and 2 show the proton NMR spectrum and carbon NMR of the obtained compound VI-1.

δ_H (CD_2Cl_2 , 400 MHz), 7.96 (d, 2H), 7.68 (dd, 2H), 7.48-7.45 (m, 6H), 7.34-7.32 (m, 4H), 7.26 (m, 2H), 7.02 (d, 2H), 6.71 (d, 4H), 4.07 (t, 4H), 1.85 (quint, 4H), 1.53 (quint, 4H), 1.44-1.35 (quint, 16H), 0.94 (t, 6H), δ_C (CD_2Cl_2 , 400 MHz), 160.0, 150.9, 143.6, 141.6, 141.19, 141.18, 140.9, 129.0, 127.6, 127.3, 127.1, 124.9, 122.7, 120.8, 115.0, 106.3, 68.8, 65.3, 32.3, 29.82, 29.81, 29.7, 26.5, 23.1, 14.3. Cald. for $C_{53}H_{56}O_2$: C, 87.80; H, 7.79. Found: C, 87.80; H, 7.77.

Example 7

Preparation of 2,7-bisbiphenyl-3'6'-bisoctyloxy-9,9'-spirobisfluorene

The compound was prepared under Suzuki reaction conditions. The compound V-1 prepared in Example 5 (0.35 mmol), 4-phenyl boric acid (0.89 mmol), tetrakis(triphenylphosphine)palladium (0.123 mmol), 2M-sodium carbonate (1.06 ml) and toluene (15 ml) were simultaneously mixed, freezed under liquid nitrogen, evacuated by a vacuum pump and then warmed to room temperature to be melted. This procedure was repeated twice, followed by injection of nitrogen. The mixed solution was refluxed for 2 days and cooled to room temperature. Thereafter, a small amount of distilled water was added, and the organic layer was filtered three

times with dichloromethane, dried with sodium sulfate. Then, the dichloromethane was concentrated under reduced pressure. Purification by silica gel chromatography using 20% dichloromethane in hexane, followed by recrystallization using a mixed solvent of hexane and dichloromethane

gave 2,7-bisbiphenyl-3',6'-bisoctyloxy-9,9'-spirobisfluorene corresponding to the compound VI-1 (Yield: 0.281 g, 90%), R_1 and R_2 in the formula of the compound VI-1 being each independently octyl and m and n being 2. The product was identified through the NMR spectrum and element analysis that it had the identical structure with the compound VI-1. FIGS. 3, 4 and 5 show the proton NMR spectrum, carbon NMR spectrum and mass spectrum of the obtained compound VI-1.

δ_H (CD_2Cl_2 , 400 MHz), 7.96 (d, 2H), 7.68 (dd, 2H), 7.59-7.52 (m, 12H), 7.40-7.38 (m, 8H), 7.01 (d, 2H), 6.68 (d, 4H), 4.04 (t, 4H), 1.80 (quint, 4H), 1.48 (quint, 4H), 1.40-1.29 (quint, 16H), 0.89 (t, 6H), δ_C (CD_2Cl_2 , 400 MHz), 159.3, 150.9, 143.5, 141.4, 140.9, 140.8, 140.2, 140.0, 129.1, 127.7, 127.5, 127.2, 126.9, 124.9, 122.5, 120.8, 114.9, 106.1, 68.7, 65.2, 32.2, 29.8, 29.7, 29.6, 26.4, 23.0, 14.2. Calc. for $C_{65}H_{64}O_2$: C, 89.00; H, 7.35. Found: C, 88.85; H, 7.68. m/z (EI-MS): Cald. 877.20 Found 877.10.

Example 8

Preparation of 4''-iodo[1,1';4',1'']terphenyl

The compound was prepared under Suzuki reaction conditions. 4,4'-diiodo-biphenyl (1.0 g, 2.46 mmol), phenyl boric acid (0.24 g, 1.970 mmol), tetrakis(triphenylphosphine)palladium (0.085 g, 0.074 mmol), 2M-sodium carbonate (12.3 ml) and toluene (76 ml) were simultaneously mixed, freezed under liquid nitrogen, evacuated by a vacuum pump and then warmed to room temperature to be melted. This procedure was repeated twice, followed by injection of nitrogen. The mixed solution was refluxed for 2 days and cooled to room temperature. Thereafter, a small amount of distilled water was added, and the organic layer was filtered three times with dichloromethane, dried with sodium sulfate. Then, the dichloromethane was concentrated under reduced pressure. Purification by silica gel chromatography using 1% dichloromethane in hexane gave

4''-iodo-[1,1';4',1'']terphenyl (Yield: 0.245 g, 35%). The structure of the product was identified through the NMR spectrum and mass spectrum. FIG. 6 shows the mass spectrum of 4''-iodo-[1,1';4',1'']terphenyl.

m/z (EI-MS): Cald. 356.01 Found 356.0

Example 9

Preparation of 2,7-diterphenyl-3'6'-bisoctyloxy-9,9'-spirobisfluorene

Bis(pinacolato)diboron (0.209 g, 0.821 mmol), 4''-iodo-[1,1';4',1'']terphenyl prepared in Example 8 (0.098 g, 0.274 mmol), [1,1'-bis(diphenylphosphino)perocine]dichloropalladium (0.007 g, 0.0082 mmol), potassium acetate (0.08 g, 0.821 mmol) and diformaldehyde (2 ml) were simultaneously mixed and evacuated by a vacuum pump, followed by injection of nitrogen and stirring at about 80°C for about 3 hours. Then, the compound V-1 prepared in Example 5 (0.10 g, 0.137 mmol), tetrakis(triphenylphosphine)palladium (0.01 g, 0.821 mmol) and 2M-sodium carbonate (0.174 g, 1.64 mmol) were mixed, freezeed under liquid nitrogen, evacuated by a vacuum pump and then warmed to room temperature to be melted. This procedure was repeated twice, followed by injection of nitrogen. The mixed solution was refluxed for 2 days and cooled to room temperature. Thereafter, a small amount of distilled water was added, and the organic layer was filtered three times with dichloromethane, dried with sodium sulfate. Then, the dichloromethane was concentrated under reduced pressure. Purification by silica gel chromatography using 1% dichloromethane in hexane gave 2,7-diterphenyl-3',6'-bisoctyloxy-9,9'-spirobifluorene (Yield: 0.085 g, 30%). The product was identified through the NMR spectrum and element analysis that it had the identical structure with the compound VI-1 of the reaction scheme 2 (R₁=octyl, R₂=octyl, m=3 and n=3). FIGS. 7 and 8 show the proton NMR spectrum and carbon NMR spectrum of the obtained compound VI-1.

δ_H (CD₂Cl₂, 400 MHz), 7.97 (d, 2H), 7.68-7.63 (m, 18H), 7.55 (d, 4H), 7.45 (m, 4H), 7.39 (m, 4H), 7.02 (t, 2H), 6.69 (d, 4H), 4.05 (t, 4H), 1.82 (quint, 4H), 1.50 (m, 4H), 1.36-1.27 (m, 16H), 0.89 (t, 6H), δ_C (CD₂Cl₂, 400 MHz), 160.0, 151.0, 143.6, 141.5, 141.0, 140.9, 140.6, 140.4, 140.2, 139.7, 139.65, 129.2, 127.77, 127.69,

127.56, 127.45, 127.26, 127.0, 124.9, 122.6, 120.9, 115.0, 106.2, 68.8, 65.3, 32.2, 29.8, 29.7, 26.5, 23.0, 14.2. m/z (EI-MS): Cald. 1029.39 Found 1029.0.

Example 10

Manufacture of EL device using dialkyl bisphenylene-spirobifluorene

compound

The EL device was manufactured using the compound VI-1 prepared in Example 7 (R_1 =octyl, R_2 =octyl, $m=2$ and $n=2$), which will now be described in detail with reference to FIG. 9. A glass substrate 10 having a transparent electrode 12 made of indium tin oxide (ITO) patterned thereon was cleaned, and then a hole-transporting layer 14, a light emitting layer 16, an electron transporting layer 18 and an anode 20 were sequentially deposited on the transparent electrode 12 by thermal deposition under a pressure of 10^{-6} torr or less.

The photoluminescence (PL) and EL spectra were measured and shown in FIG. 10. In FIG. 10, the maximum emission peak is 402 nm and blue EL is displayed in the range from 400 to 500 nm. The CIE color coordinate of light emitted is (0.16, 0.09), that is, blue with high color purity, exhibiting high luminance and high emission efficiency.

While the invention has been described in terms of specific examples, the substrate 10 may be formed of not only glass but also polycarbonate, polyimide, polyethyleneterephthalate or polyethylenenaphthalate. Also, the transparent electrode 12 may be formed of indium tin oxide, tin oxide, zinc oxide or combinations thereof. The transparent electrode 12 may be formed by mixing these exemplary materials in an appropriate ratio according to the requirements in view of transmissivity and conductivity, using a sputtering method.

The bisphenylene-spirobifluorene compound according to the present invention is a monomer having an alkyl substituent introduced to a new position. The alkyl side chains can serve to reduce an intermolecular interaction by preventing π - π interactions of phenylene groups of spirobifluorene, resulting in reducing the excimer formation and keeping the color purity. In other words, in the spirobifluorene compound according to the present invention, one fluorene ring has an alkyl

substituent for increasing solubility in organic solvents and reducing intermolecular interaction, and two phenylene groups are introduced to the other fluorene ring for enhancing blue EL property. Thus, the bisphenylene-spirobifluorene compound according to the present invention has high solubility in organic solvents as well as excellent color purity and color stability, thereby having high processability. Also, the present invention can provide an organic blue EL material having properties good enough to meet requirements as electronic materials, including transmissivity, environment resistance, adhesive strength with respect to a substrate, film-forming property, electric field stability and so on. Formation of a light emitting layer of an EL device using the compound according to the present invention noticeably reduces crystallization typically occurring to planar organic materials during formation of a thin film for forming the light emitting layer, thereby extending the life span of the EL device.

While this invention has been particularly shown and described with reference to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention as defined by the appended claims.